

2장 참고문헌

Ackert CL, Gittens JE, O'Brien MJ, Eppig JJ, Kidder GM. Intercellular communication via connexin43 gap junctions is required for ovarian folliculogenesis in the mouse. *Dev Biol* 2001; 233: 258-70.

Alak BM, Smith GD, Woodruff TK, Stouffer RL, Wolf DP. Enhancement of primate oocyte maturation and fertilization in vitro by inhibin A and activin A. *Fertil Steril* 1996; 66: 646-53.

Anasti JN, Kalantaridou SN, Kimzey LM, George M, Nelson LM. Human follicle fluid vascular endothelial growth factor concentrations are correlated with luteinization in spontaneously developing follicles. *Hum Reprod* 1998; 13: 1144-7.

Andersen CY. Characteristics of human follicular fluid associated with successful conception after in vitro fertilization. *J Clin Endocrinol Metab* 1993; 77: 1227-34.

Andersen CY, Byskov AG. Estradiol and regulation of anti-müllerian hormone, inhibin-A, and inhibin-B secretion: analysis of small antral and preovulatory human follicles' fluid. *J Clin Endocrinol Metab* 2006; 91: 4064-9.

Asakura H, Zwain IH, Yen SS. Expression of genes encoding corticotropin-releasing factor (CRF), type 1 CRF receptor, and CRF-binding protein and localization of the gene products in the human ovary. *J Clin Endocrinol Metab* 1997; 82: 2720-5.

Auletta FJ, Flint AP. Mechanisms controlling corpus luteum function in sheep, cows, nonhuman primates, and women especially in relation to the time of luteolysis. *Endocr Rev* 1988; 9: 88-105.

Azhar S, Tsai L, Medicherla S, Chandrasekhar Y, Giudice L, Reaven E. Human granulosa cells use high density lipoprotein cholesterol for steroidogenesis. *J Clin Endocrinol Metab* 1998; 83: 983-91.

Baker TG. A quantitative and cytological study of germ cells in human ovaries. *Proc Roy Soc Lond* 1963; 158: 417-33.

Baker TG, Scrimgeour JB. Development of the gonad in normal and anencephalic human fetuses. *J Reprod Fertil* 1980; 60: 193-9.

Bassett SG, Little-Ihrig LL, Mason JI, Zeleznik AJ. Expression of messenger ribonucleic acids that encode for 3 beta-hydroxysteroid dehydrogenase and cholesterol side-chain cleavage enzyme throughout the luteal phase of the macaque menstrual cycle. *J Clin Endocrinol Metab* 1991; 72: 362-6.

Barnes RB, Namnoum AB, Rosenfield RL, Layman LC. The role of LH and FSH in ovarian androgen secretion and ovarian follicular development: clinical studies in a patient with isolated FSH deficiency and multicystic ovaries. *Hum Reprod* 2002; 17: 88-91.

Barreca A, Artini PG, Del Monte P, Ponzani P, Pasquini P, Cariola G, et al. In vivo and in vitro effect of growth hormone on estradiol secretion by human granulosa cells. *J Clin Endocrinol Metab* 1993; 77: 61-7.

Ben-Ami I, Armon L, Freimann S, Strassburger D, Ron-El R, Amsterdam A. EGF-like growth factors as LH mediators in the human corpus luteum. *Hum Reprod* 2009; 24: 176-84.

Ben-Ami I, Freimann S, Armon L, Dantes A, Strassburger D, Friedler S, et al. PGE2 up-regulates EGF-like growth factor



biosynthesis in human granulosa cells: new insights into the coordination between PGE2 and LH in ovulation. *Mol Hum Reprod* 2006; 12: 593-9.

Ben-Chetrit A, Gotlieb L, Wong PY, Casper RF. Ovarian response to recombinant human follicle-stimulating hormone in luteinizing hormone-depleted women: examination of the two cell, two gonadotropin theory. *Fertil Steril* 1996; 65: 711-7.

Besecke LM, Guendner MJ, Sluss PA, Polak AG, Woodruff TK, Jameson JL, et al. Pituitary follistatin regulates activin-mediated production of follicle-stimulating hormone during the rat estrous cycle. *Endocrinology* 1997; 138: 2841-8.

Bergh C, Carlsson B, Olsson JH, Selleskog U, Hillensjö T. Regulation of androgen production in cultured human thecal cells by insulin-like growth factor I and insulin. *Fertil Steril* 1993; 59: 323-31.

Bilezikjian LM, Corrigan AZ, Blount AL, Vale WW. Pituitary follistatin and inhibin subunit messenger ribonucleic acid levels are differentially regulated by local and hormonal factors. *Endocrinology* 1996; 137: 4277-84.

Bogan RL, Murphy MJ, Stouffer RL, Hennebold JD. Prostaglandin synthesis, metabolism, and signaling potential in the rhesus macaque corpus luteum throughout the luteal phase of the menstrual cycle. *Endocrinology* 2008; 149: 5861-71.

Braden TD, Conn PM. Activin-A stimulates the synthesis of gonadotropin-releasing hormone receptors. *Endocrinology* 1992; 130: 2101-5.

Brannian JD, Stouffer RL. Progesterone production by monkey luteal cell subpopulations at different stages of the menstrual cycle: changes in agonist responsiveness. *Biol Reprod* 1991; 44: 141-9.

Buckler HM, Healy DL, Burger HG. Purified FSH stimulates production of inhibin by the human ovary. *J Endocrinol* 1989; 122: 279-85.

Cahill DJ, Wardle PG, Harlow CR, Hull MG. Onset of the preovulatory luteinizing hormone surge: diurnal timing and critical follicular prerequisites. *Fertil Steril* 1998; 70:56-9.

Castro A, Castro O, Troncoso JL, Kohen P, Simón C, Vega M, et al. Luteal leukocytes are modulators of the steroidogenic process of human mid-luteal cells. *Hum Reprod* 1998; 13: 1584-9.

Cataldo NA, Giudice LC. Insulin-like growth factor binding protein profiles in human ovarian follicular fluid correlate with follicular functional status. *J Clin Endocrinol Metab* 1992; 74: 821-9.

Chabab A, Hedon B, Arnal F, Diafouka F, Bressot N, Flandre O, et al. Follicular steroids in relation to oocyte development and human ovarian stimulation protocols. *Hum Reprod* 1986; 1: 449-54.

Chaffin CL, Stouffer RL. Expression of matrix metalloproteinases and their tissue inhibitor messenger ribonucleic acids in macaque periovulatory granulosa cells: time course and steroid regulation. *Biol Reprod* 1999; 61: 14-21.

Chandrasekher YA, Van Dessel HJ, Fauser BC, Giudice LC. Estrogen- but not androgen-dominant human ovarian follicular fluid contains an insulin-like growth factor binding protein-4 protease. *J Clin Endocrinol Metab* 1995; 80: 2734-9.

Chang RJ, Gougeon A, Erickson GF. Evidence for a neutrophil-interleukin-8 system in human folliculogenesis. *Am J Obstet Gynecol* 1998; 178: 650-7.

Chappel SC, Resko JA, Norman RL, Spies HG. Studies in rhesus monkeys on the site where estrogen inhibits gonadotropins: delivery of 17 beta-estradiol to the hypothalamus and pituitary gland. *J Clin Endocrinol Metab* 1981; 52: 1-8.

Chikazawa K, Araki S, Tamada T. Morphological and endocrinological studies on follicular development during the human menstrual cycle. *J Clin Endocrinol Metab* 1986; 62:305-13.

Clark JR, Dierschke DJ, Wolf RC. Hormonal regulation of ovarian folliculogenesis in rhesus monkeys: III. Atresia of the preovulatory follicle induced by exogenous steroids and subsequent follicular development. *Biol Reprod* 1981; 25: 332-41.

Collins RL, Hodgen GD. Blockade of the spontaneous midcycle gonadotropin surge in monkeys by RU 486: a progesterone antagonist or agonist? *J Clin Endocrinol Metab* 1986; 63: 1270-6.

Conneely OM, Mulac-Jericevic B, Lydon JP, De Mayo FJ. Reproductive functions of the progesterone receptor isoforms: lessons from knock-out mice. *Mol Cell Endocrinol* 2001; 179: 97-103.

Couzinet B, Brailly S, Bouchard P, Schaison G. Progesterone stimulates luteinizing hormone secretion by acting directly on the pituitary. *J Clin Endocrinol Metab* 1992; 74: 374-8.

de Koning J. Gonadotrophin surge-inhibiting/attenuating factor governs luteinizing hormone secretion during the ovarian cycle: physiology and pathology. *Hum Reprod* 1995; 10: 2854-61.

Demura R, Suzuki T, Tajima S, Mitsuhashi S, Odagiri E, Demura H, et al. Human plasma free activin and inhibin levels during the menstrual cycle. *J Clin Endocrinol Metab* 1993; 76: 1080-2.

Diaz FJ, Sugiura K, Eppig JJ. Regulation of Pcsk6 expression during the preantral to antral follicle transition in mice: opposing roles of FSH and oocytes. *Biol Reprod* 2008; 78: 176-83.

Diaz FJ, Wigglesworth K, Eppig JJ. Oocytes are required for the preantral granulosa cell to cumulus cell transition in mice. *Dev Biol* 2007; 305: 300-11.

Di Blasio AM, Viganó P, Ferrari A. Insulin-like growth factor-II stimulates human granulosa-luteal cell proliferation in vitro. *Fertil Steril* 1994; 61: 483-7.

Dissen GA, Garcia-Rudaz C, Ojeda SR. Role of neurotrophic factors in early ovarian development. *Semin Reprod Med* 2009; 27: 24-31.

Dor J, Ben-Shlomo I, Lunenfeld B, Pariente C, Levran D, Karasik A, et al. Insulin-like growth factor-I (IGF-I) may not be essential for ovarian follicular development: evidence from IGF-I deficiency. *J Clin Endocrinol Metab* 1992; 74: 539-42.

Dodson WC, Hughes CL, Whitesides DB, Haney AF. The effect of leuprolide acetate on ovulation induction with human menopausal gonadotropins in polycystic ovary syndrome. *J Clin Endocrinol Metab* 1987; 65: 95-100.

Duffy DM, Dozier BL, Seachord CL. Prostaglandin dehydrogenase and prostaglandin levels in periovulatory follicles: implications for control of primate ovulation by prostaglandin E2. *J Clin Endocrinol Metab* 2005; 90: 1021-7.

Duncan WC, McNeilly AS, Illingworth PJ. The effect of luteal "rescue" on the expression and localization of matrix metalloproteinases and their tissue inhibitors in the human corpus luteum. *J Clin Endocrinol Metab* 1998; 83: 2470-8.



Durlinger AL, Grujters MJ, Kramer P, Karels B, Ingraham HA, Nachtigal MW, et al. Anti-Müllerian hormone inhibits initiation of primordial follicle growth in the mouse ovary. *Endocrinology* 2002; 143: 1076-84.

Enmark E, Peltö-Huikko M, Grandien K, Lagercrantz S, Lagercrantz J, Fried G, et al. Human estrogen receptor beta-gene structure, chromosomal localization, and expression pattern. *J Clin Endocrinol Metab* 1997; 82: 4258-65.

Erickson GF, Magoffin DA, Dyer CA, Hofeditz C. The ovarian androgen producing cells: a review of structure/function relationships. *Endocr Rev* 1985; 6: 371-99.

Erickson GF. An analysis of follicle development and ovum maturation. *Seminars Reprod Endocrinol* 1986; 4: 233.

Erden HF, Zwain IH, Asakura H, Yen SS. Corticotropin-releasing factor inhibits luteinizing hormone-stimulated P450c17 gene expression and androgen production by isolated thecal cells of human ovarian follicles. *J Clin Endocrinol Metab* 1998; 83: 448-52.

Eppig JJ, Chesnel F, Hirao Y, O'Brien MJ, Pendola FL, Watanabe S, et al. Oocyte control of granulosa cell development: how and why. *Hum Reprod* 1997; 12: 127-32.

Evans WS, Sollenberger MJ, Booth RA Jr, Rogol AD, Urban RJ, Carlsen EC, et al. Contemporary aspects of discrete peak-detection algorithms. II. The paradigm of the luteinizing hormone pulse signal in women. *Endocr Rev* 1992; 13: 81-104.

Filicori M, Cognigni GE, Tabarelli C, Pocognoli P, Taraborrelli S, Spetoli D, et al. Stimulation and growth of antral ovarian follicles by selective LH activity administration in women. *J Clin Endocrinol Metab* 2002; 87: 1156-61.

Filicori M, Cognigni GE, Ciampaglia W. Effects of LH on oocyte yield and developmental competence. *Hum Reprod* 2003; 18: 1357-8.

Filicori M, Butler JP, Crowley WF Jr. Neuroendocrine regulation of the corpus luteum in the human. Evidence for pulsatile progesterone secretion. *J Clin Invest* 1984; 73: 1638-47.

Fletcher WH, Greenan JR. Receptor mediated action without receptor occupancy. *Endocrinology* 1985; 116: 1660-2.

Fridén BE, Runesson E, Hahlin M, Brännström M. Evidence for nitric oxide acting as a luteolytic factor in the human corpus luteum. *Mol Hum Reprod* 2000; 6: 397-403.

Fukuda M, Fukuda K, Andersen CY, Byskov AG. Right-sided ovulation favours pregnancy more than left-sided ovulation. *Hum Reprod* 2000; 15: 1921-6.

Fukuda M, Fukuda K, Andersen CY, Byskov AG. Characteristics of human ovulation in natural cycles correlated with age and achievement of pregnancy. *Hum Reprod* 2001; 16: 2501-7.

Fritz MA, McLachlan RI, Cohen NL, Dahl KD, Bremner WJ, Soules MR. Onset and characteristics of the midcycle surge in bioactive and immunoactive luteinizing hormone secretion in normal women: influence of physiological variations in periovulatory ovarian steroid hormone secretion. *J Clin Endocrinol Metab* 1992; 75: 489-93.

Fritz MA, Speroff L. Regulation of the Menstrual Cycle In: *Clinical Gynecologic Endocrinology and Infertility*. 8th ed. Lippincott Williams & Wilkins 2011; p.201.

- Giudice LC. Insulin-like growth factors and ovarian follicular development. *Endocr Rev* 1992; 13: 641-69.
- Girsh E, Milvae RA, Wang W, Meidan R. Effect of endothelin-1 on bovine luteal cell function: role in prostaglandin F₂α-induced antisteroidogenic action. *Endocrinology* 1996; 137: 1306-12.
- Golos TG, Soto EA, Tureck RW, Strauss JF 3rd. Human chorionic gonadotropin and 8-bromo-adenosine 3',5'-monophosphate stimulate [125I]low density lipoprotein uptake and metabolism by luteinized human granulosa cells in culture. *J Clin Endocrinol Metab* 1985; 61: 633-8.
- Goodman AL, Hodgen GD. The ovarian triad of the primate menstrual cycle. *Recent Prog Horm Res* 1983; 39: 1-73.
- Gougeon A. Dynamics of follicular growth in the human: a model from preliminary results. *Hum Reprod* 1986; 1:81-7.
- Gougeon A. Regulation of ovarian follicular development in primates: facts and hypotheses. *Endocr Rev* 1996; 17: 121-55.
- Gore BZ, Caldwell BV, Speroff L. Estrogen-induced human luteolysis. *J Clin Endocrinol Metab* 1973; 36: 615-7.
- Granot I, Dekel N. Developmental expression and regulation of the gap junction protein and transcript in rat ovaries. *Mol Reprod Dev* 1997; 47:231-9.
- Granot I, Dekel N. The ovarian gap junction protein connexin43: regulation by gonadotropins. *Trends Endocrinol Metab* 2002; 13: 310-3.
- Grazul-Bilska AT, Redmer DA, Reynolds LP. Effects of luteinizing hormone and prostaglandin F(2α) on gap junctional intercellular communication of ovine luteal cells throughout the estrous cycle. *Endocrine* 1996; 5: 225-33.
- Groome NP, Illingworth PJ, O'Brien M, Pai R, Rodger FE, Mather JP, et al. Measurement of dimeric inhibin B throughout the human menstrual cycle. *J Clin Endocrinol Metab* 1996; 81: 1401-5.
- Grimes RW, Samaras SE, Barber JA, Shimasaki S, Ling N, Hammond JM. Gonadotropin and cAMP modulation of IGE binding protein production in ovarian granulosa cells. *Am J Physiol* 1992; 262: E497-503.
- Hall JE, Taylor AE, Martin KA, Rivier J, Schoenfeld DA, Crowley WF Jr. Decreased release of gonadotropin-releasing hormone during the preovulatory midcycle luteinizing hormone surge in normal women. *Proc Natl Acad Sci U S A* 1994; 91: 6894-8.
- Hedricks C, Piccinino LJ, Udry JR, Chimbira TH. Peak coital rate coincides with onset of luteinizing hormone surge. *Fertil Steril* 1987; 48: 234-8.
- Hernandez ER, Hurwitz A, Payne DW, Dharmarajan AM, Purchio AF, Adashi EY. Transforming growth factor-beta 1 inhibits ovarian androgen production: gene expression, cellular localization, mechanisms(s), and site(s) of action. *Endocrinology* 1990; 127: 2804-11.
- Hibbert ML, Stouffer RL, Wolf DP, Zelinski-Wooten MB. Midcycle administration of a progesterone synthesis inhibitor prevents ovulation in primates. *Proc Natl Acad Sci U S A* 1996; 93: 1897-901.
- Hild-Petito S, Stouffer RL, Brenner RM. Immunocytochemical localization of estradiol and progesterone receptors in the monkey ovary throughout the menstrual cycle. *Endocrinology* 1988; 123: 2896-905.



Hillier SG, Yong EL, Illingworth PJ, Baird DT, Schwall RH, Mason AJ. Effect of recombinant activin on androgen synthesis in cultured human thecal cells. *J Clin Endocrinol Metab* 1991; 72: 1206-11.

Hoff JD, Quigley ME, Yen SS. Hormonal dynamics at midcycle: a reevaluation. *J Clin Endocrinol Metab* 1983; 57: 792-6.

Itskovitz J, Sealey JE. Ovarian prorenin-renin-angiotensin system. *Obstet Gynecol Surv* 1987; 42: 545-51.

Jia XC, Kessel B, Welsh TH Jr, Hsueh AJ. Androgen inhibition of follicle-stimulating hormone-stimulated luteinizing hormone receptor formation in cultured rat granulosa cells. *Endocrinology* 1985; 117: 13-22.

Judd S, Terry A, Petrucco M, White G. The source of pulsatile secretion of progesterone during the human follicular phase. *J Clin Endocrinol Metab* 1992; 74: 299-305.

Juengel JL, McNatty KP. The role of proteins of the transforming growth factor-beta superfamily in the intraovarian regulation of follicular development. *Hum Reprod Update* 2005; 11: 143-60.

Kang SK, Tai CJ, Nathwani PS, Leung PC. Differential regulation of two forms of gonadotropin-releasing hormone messenger ribonucleic acid in human granulosa-luteal cells. *Endocrinology* 2001; 142: 182-92.

Katt JA, Duncan JA, Herbon L, Barkan A, Marshall JC. The frequency of gonadotropin-releasing hormone stimulation determines the number of pituitary gonadotropin-releasing hormone receptors. *Endocrinology* 1985; 116: 2113-5.

Kettel LM, DePaolo LV, Morales AJ, Apter D, Ling N, Yen SS. Circulating levels of follistatin from puberty to menopause. *Fertil Steril* 1996; 65: 472-6.

Kim JH, Seibel MM, MacLaughlin DT, Donahoe PK, Ransil BJ, Hametz PA, et al. The inhibitory effects of müllerian-inhibiting substance on epidermal growth factor induced proliferation and progesterone production of human granulosa-luteal cells. *J Clin Endocrinol Metab* 1992; 75: 911-7.

Kitaoka M, Kojima I, Ogata E. Activin-A: a modulator of multiple types of anterior pituitary cells. *Biochem Biophys Res Commun* 1988; 157: 48-54.

Klein NA, Illingworth PJ, Groome NP, McNeilly AS, Battaglia DE, Soules MR. Decreased inhibin B secretion is associated with the monotropic FSH rise in older, ovulatory women: a study of serum and follicular fluid levels of dimeric inhibin A and B in spontaneous menstrual cycles. *J Clin Endocrinol Metab* 1996; 81: 2742-5.

Knight PG, Glister C. TGF-beta superfamily members and ovarian follicle development. *Reproduction* 2006; 132: 191-206.

Kobayashi M, Nakano R, Ooshima A. Immunohistochemical localization of pituitary gonadotrophins and gonadal steroids confirms the 'two-cell, two-gonadotrophin' hypothesis of steroidogenesis in the human ovary. *J Endocrinol* 1990; 126: 483-8.

Kokia E, Hurwitz A, Ricciarelli E, Tedeschi C, Resnick CE, Mitchell MD, et al. Interleukin-1 stimulates ovarian prostaglandin biosynthesis: evidence for heterologous contact-independent cell-cell interaction. *Endocrinology* 1992; 130: 3095-7.

Lau AL, Kumar TR, Nishimori K, Bonadio J, Matzuk MM. Activin betaC and betaE genes are not essential for mouse liver growth, differentiation, and regeneration. *Mol Cell Biol* 2000; 20: 6127-37.

Lee A, Christenson LK, Patton PE, Burry KA, Stouffer RL. Vascular endothelial growth factor production by human luteinized granulosa cells in vitro. *Hum Reprod* 1997; 12: 2756-61.

Lei ZM, Chegini N, Rao CV. Quantitative cell composition of human and bovine corpora lutea from various reproductive states. *Biol Reprod* 1991; 44: 1148-56.

Lenton EA, Landgren BM, Sexton L. Normal variation in the length of the luteal phase of the menstrual cycle: identification of the short luteal phase. *Br J Obstet Gynaecol* 1984; 91: 685-9.

Li W, Yuen BH, Leung PC. Inhibition of progesterin accumulation by activin-A in human granulosa cells. *J Clin Endocrinol Metab* 1992; 75: 285-9.

Mais V, Kazer RR, Cetel NS, Rivier J, Vale W, Yen SS. The dependency of folliculogenesis and corpus luteum function on pulsatile gonadotropin secretion in cycling women using a gonadotropin-releasing hormone antagonist as a probe. *J Clin Endocrinol Metab* 1986; 62: 1250-5.

Mason HD, Cwyfan-Hughes SC, Heinrich G, Franks S, Holly JM. Insulin-like growth factor (IGF) I and II, IGF-binding proteins, and IGF-binding protein proteases are produced by theca and stroma of normal and polycystic human ovaries. *J Clin Endocrinol Metab* 1996; 81: 276-84.

Maas S, Jarry H, Teichmann A, Rath W, Kuhn W, Wuttke W. Paracrine actions of oxytocin, prostaglandin F2 alpha, and estradiol within the human corpus luteum. *J Clin Endocrinol Metab* 1992; 74: 306-12.

Markosyan N, Duffy DM. Prostaglandin E2 acts via multiple receptors to regulate plasminogen-dependent proteolysis in the primate periovulatory follicle. *Endocrinology* 2009; 150: 435-44.

McLachlan RI, Cohen NL, Vale WW, Rivier JE, Burger HG, Bremner WJ, et al. The importance of luteinizing hormone in the control of inhibin and progesterone secretion by the human corpus luteum. *J Clin Endocrinol Metab* 1989; 68: 1078-85.

McNatty KP, Makris A, DeGrazia C, Osathanondh R, Ryan KJ. The production of progesterone, androgens, and estrogens by granulosa cells, thecal tissue, and stromal tissue from human ovaries in vitro. *J Clin Endocrinol Metab* 1979; 49: 687-99.

McNatty KP, Makris A, De Grazia C, Osathanondh R, Ryan KJ. Steroidogenesis by recombined follicular cells from the human ovary in vitro. *J Clin Endocrinol Metab* 1980; 51: 1286-92.

McNeely MJ, Soules MR. The diagnosis of luteal phase deficiency: a critical review. *Fertil Steril* 1988; 50: 1-15.

Meyer AC, Papadimitriou JC, Silverberg SG, Sharara FI. Secondary amenorrhea and infertility caused by an inhibin-B-producing ovarian fibrothecoma. *Fertil Steril* 2000; 73: 258-60.

Miceli F, Minici F, Garcia Pardo M, Navarra P, Proto C, Mancuso S, et al. Endothelins enhance prostaglandin (PGE(2) and PGF(2alpha)) biosynthesis and release by human luteal cells: evidence of a new paracrine/autocrine regulation of luteal function. *J Clin Endocrinol Metab* 2001; 86: 811-7.

Montgomery Rice V, Limback SD, Roby KF, Terranova PF. Differential responses of granulosa cells from small and large follicles to follicle stimulating hormone (FSH) during the menstrual cycle and acyclicity: effects of tumour necrosis factor-alpha. *Hum Reprod* 1998; 13: 1285-91.



Myers M, Gay E, McNeilly AS, Fraser HM, Duncan WC. In vitro evidence suggests activin-A may promote tissue remodeling associated with human luteolysis. *Endocrinology* 2007; 148: 3730-9.

O' Sullivan MJ, Stamouli A, Thomas EJ, Richardson MC. Gonadotrophin regulation of production of tissue inhibitor of metalloproteinases-1 by luteinized human granulosa cells: a potential mechanism for luteal rescue. *Mol Hum Reprod* 1997; 3: 405-10.

Ottowitz WE, Dougherty DD, Fischman AJ, Hall JE. [18F]2-fluoro-2-deoxy-D-glucose positron emission tomography demonstration of estrogen negative and positive feedback on luteinizing hormone secretion in women. *J Clin Endocrinol Metab* 2008; 93: 3208-14.

Panigone S, Hsieh M, Fu M, Persani L, Conti M. Luteinizing hormone signaling in preovulatory follicles involves early activation of the epidermal growth factor receptor pathway. *Mol Endocrinol* 2008; 22: 924-36.

Pauerstein CJ, Eddy CA, Croxatto HD, Hess R, Siler-Khodr TM, Croxatto HB. Temporal relationships of estrogen, progesterone, and luteinizing hormone levels to ovulation in women and infrahuman primates. *Am J Obstet Gynecol* 1978; 130: 876-86.

Peluffo MC, Young KA, Hennebold JD, Stouffer RL. Expression and regulation of tumor necrosis factor (TNF) and TNF-receptor family members in the macaque corpus luteum during the menstrual cycle. *Mol Reprod Dev* 2009; 76: 367-78.

Peng XR, Hsueh AJ, Ny T. Transient and cell-specific expression of tissue-type plasminogen activator and plasminogen-activator-inhibitor type 1 results in controlled and directed proteolysis during gonadotropin-induced ovulation. *Eur J Biochem* 1993; 214: 147-56.

Piquette GN, Simón C, el Danasouri I, Frances A, Polan ML. Gene regulation of interleukin-1 beta, interleukin-1 receptor type I, and plasminogen activator inhibitor-1 and -2 in human granulosa-luteal cells. *Fertil Steril*. 1994; 62: 760-70.

Retamales I, Carrasco I, Troncoso JL, Las Heras J, Devoto L, Vega M. Morpho-functional study of human luteal cell subpopulations. *Hum Reprod* 1994; 9: 591-6.

Richards JS, Jahnsen T, Hedin L, Lifka J, Ratoosh S, Durica JM, et al. Ovarian follicular development: from physiology to molecular biology. *Recent Prog Horm Res* 1987; 43: 231-76.

Richards JS, Liu Z, Shimada M. Immune-like mechanisms in ovulation. *Trends Endocrinol Metab* 2008; 19: 191-6.

Robertson DM. Follistatin/activin-binding protein. *Trends Endocrinol Metab* 1992; 3: 65-8.

Sasano H, Okamoto M, Mason JI, Simpson ER, Mendelson CR, Sasano N, et al. Immunolocalization of aromatase, 17 alpha-hydroxylase and side-chain-cleavage cytochromes P-450 in the human ovary. *J Reprod Fertil* 1989; 85: 163-9.

Schipper I, Hop WC, Fauser BC. The follicle-stimulating hormone (FSH) threshold/window concept examined by different interventions with exogenous FSH during the follicular phase of the normal menstrual cycle: duration, rather than magnitude, of FSH increase affects follicle development. *J Clin Endocrinol Metab* 1998; 83: 1292-8.

Sela-Abramovich S, Galiani D, Nevo N, Dekel N. Inhibition of rat oocyte maturation and ovulation by nitric oxide: mechanism of action. *Biol Reprod* 2008; 78: 1111-8.

Shimasaki S, Ling N. Identification and molecular characterization of insulin-like growth factor binding proteins (IGFBP-1, -2, -3, -4, -5 and -6). *Prog Growth Factor Res* 1991; 3: 243-66.

Smith G, Roberts R, Hall C, Nuki G. Reversible ovulatory failure associated with the development of luteinized unruptured follicles in women with inflammatory arthritis taking non-steroidal anti-inflammatory drugs. *Br J Rheumatol* 1996; 35: 458-62.

Streuli I, Fraise T, Pillet C, Ibecheole V, Bischof P, de Ziegler D. Serum antimüllerian hormone levels remain stable throughout the menstrual cycle and after oral or vaginal administration of synthetic sex steroids. *Fertil Steril* 2008; 90: 395-400.

Suzuki T, Sasano H, Takaya R, Fukaya T, Yajima A, Nagura H. Cyclic changes of vasculature and vascular phenotypes in normal human ovaries. *Hum Reprod* 1998; 13: 953-9.

Taylor PD, Wilson H, Hillier SG, Wiegand SJ, Fraser HM. Effects of inhibition of vascular endothelial growth factor at time of selection on follicular angiogenesis, expansion, development and atresia in the marmoset. *Mol Hum Reprod* 2007; 13: 729-36.

Tedeschi C, Hazum E, Kokia E, Ricciarelli E, Adashi EY, Payne DW. Endothelin-1 as a luteinization inhibitor: inhibition of rat granulosa cell progesterone accumulation via selective modulation of key steroidogenic steps affecting both progesterone formation and degradation. *Endocrinology* 1992; 131: 2476-8.

Thierry van Dessel HJ, Chandrasekher Y, Yap OW, Lee PD, Hintz RL, Faessen GH, et al. Serum and follicular fluid levels of insulin-like growth factor I (IGF-I), IGF-II, and IGF-binding protein-1 and -3 during the normal menstrual cycle. *J Clin Endocrinol Metab* 1996; 81: 1224-31.

Tilly JL, Kowalski KI, Schomberg DW, Hsueh AJ. Apoptosis in atretic ovarian follicles is associated with selective decreases in messenger ribonucleic acid transcripts for gonadotropin receptors and cytochrome P450 aromatase. *Endocrinology* 1992; 131: 1670-6.

Trombly DJ, Woodruff TK, Mayo KE. Roles for transforming growth factor beta superfamily proteins in early folliculogenesis. *Semin Reprod Med* 2009; 27: 14-23.

Van Deerlin PG, Cekleniak N, Coutifaris C, Boyd J, Strauss JF 3rd. Evidence for the oligoclonal origin of the granulosa cell population of the mature human follicle. *J Clin Endocrinol Metab* 1997; 82: 3019-24.

Vanderhyden BC, Telfer EE, Eppig JJ. Mouse oocytes promote proliferation of granulosa cells from preantral and antral follicles in vitro. *Biol Reprod* 1992; 46: 1196-204.

Vande Wiele RL, Bogumil J, Dyrenfurth I, Ferin M, Jewelewicz R, Warren M, et al. Mechanisms regulating the menstrual cycle in women. *Recent Prog Horm Res* 1970; 26: 63-103.

van Rooij IA, Broekmans FJ, Scheffer GJ, Looman CW, Habbema JD, de Jong FH, et al. Serum antimüllerian hormone levels best reflect the reproductive decline with age in normal women with proven fertility: a longitudinal study. *Fertil Steril* 2005; 83: 979-87.

Voutilainen R, Franks S, Mason HD, Martikainen H. Expression of insulin-like growth factor (IGF), IGF-binding protein, and IGF receptor messenger ribonucleic acids in normal and polycystic ovaries. *J Clin Endocrinol Metab* 1996; 81: 1003-8.

Weenen C, Laven JS, Von Berg AR, Cranfield M, Groome NP, Visser JA, et al. Anti-Müllerian hormone expression pattern



in the human ovary: potential implications for initial and cyclic follicle recruitment, *Mol Hum Reprod* 2004; 10: 77-83.

Welt CK, Martin KA, Taylor AE, Lambert-Messerlian GM, Crowley WF Jr, Smith JA, et al. Frequency modulation of follicle-stimulating hormone (FSH) during the luteal-follicular transition: evidence for FSH control of inhibin B in normal women, *J Clin Endocrinol Metab* 1997; 82: 2645-52.

Wildt L, Hutchison JS, Marshall G, Pohl CR, Knobil E. On the site of action of progesterone in the blockade of the estradiol-induced gonadotropin discharge in the rhesus monkey. *Endocrinology* 1981; 109: 1293-4.

Wulff C, Wilson H, Largue P, Duncan WC, Armstrong DG, Fraser HM. Angiogenesis in the human corpus luteum: localization and changes in angiopoietins, tie-2, and vascular endothelial growth factor messenger ribonucleic acid, *J Clin Endocrinol Metab* 2000; 85: 4302-9.

Wunder DM, Bersinger NA, Yared M, Kretschmer R, Birkhäuser MH. Statistically significant changes of antimüllerian hormone and inhibin levels during the physiologic menstrual cycle in reproductive age women, *Fertil Steril* 2008; 89: 927-33.

Xiao S, Robertson DM, Findlay JK. Effects of activin and follicle-stimulating hormone (FSH)-suppressing protein/follistatin on FSH receptors and differentiation of cultured rat granulosa cells, *Endocrinology* 1992; 131: 1009-16.

Yong EL, Baird DT, Hillier SG. Mediation of gonadotrophin-stimulated growth and differentiation of human granulosa cells by adenosine-3',5'-monophosphate: one molecule, two messages. *Clin Endocrinol (Oxf)* 1992; 37: 51-8.

Yoshimura Y, Santulli R, Atlas SJ, Fujii S, Wallach EE. The effects of proteolytic enzymes on in vitro ovulation in the rabbit, *Am J Obstet Gynecol* 1987; 157: 468-75.

Zelevnik AJ, Schuler HM, Reichert LE Jr. Gonadotropin-binding sites in the rhesus monkey ovary: role of the vasculature in the selective distribution of human chorionic gonadotropin to the preovulatory follicle. *Endocrinology* 1981; 109: 356-62.

Zelinski-Wooten MB, Hutchison JS, Hess DL, Wolf DP, Stouffer RL. Follicle stimulating hormone alone supports follicle growth and oocyte development in gonadotrophin-releasing hormone antagonist-treated monkeys. *Hum Reprod* 1995; 10: 1658-66.