

 참고 문헌

1. Abou-Setta AM, Al-Inany HG, Mansour RT, Serour GI, Aboulghar MA. Soft versus firm embryo transfer catheters for assisted reproduction: a systematic review and meta-analysis. *Hum Reprod* 2005;20:3114-21.
2. Bennett SJ, Waterstone JJ, Cheng WC, Parsons J. Complications of transvaginal ultrasound-directed follicle aspiration: a review of 2670 consecutive procedures. *J Assist Reprod Genet* 1993;10:72-7.
3. Brown J, Buckingham K, Abou-Setta AM, Buckett W. Ultrasound versus 'clinical touch' for catheter guidance during embryo transfer in women. *Cochrane Database Syst Rev* 2010;1:CD006107.
4. Chang YS, Jun JK, Choi YM, Moon SY. The present status of assisted reproductive technology in Asia-Oceania. *Asia Oceania J Obstet Gynaecol* 1994;20:331-44.
5. Dorn C, Reinsberg J, Schlebusch H, Prietl G, van der Ven H, Krebs D. Serum oxytocin concentration during embryo transfer procedure. *Eur J Obstet Gynecol Reprod Biol* 1999;87:77-80.
6. Fanchin R, Righini C, Olivennes F, Taylor S, de Ziegler D, Frydman R. Uterine contractions at the time of embryo transfer alter pregnancy rates after in-vitro fertilization. *Hum Reprod* 1998;13:1968-74.
7. Frankfurter D, Trimarchi JB, Silva CP, Keefe DL. Middle to lower uterine segment embryo transfer improves implantation and pregnancy rates compared with fundal embryo transfer. *Fertil Steril* 2004;81:1273-7.
8. Hannoun A, Awwad J, Zreik T, Ghaziri G, Abu-Musa A. Effect of betadine vaginal preparation during oocyte aspiration in in vitro fertilization cycles on pregnancy outcome. *Gynecol Obstet Invest* 2008;66:274-8.
9. Haydardedeoglu B, Cok T, Kilicdag EB, Parlakgumus AH, Simsek E, Bagis T. In vitro fertilization-intracytoplasmic sperm injection outcomes in single- versus double-lumen oocyte retrieval needles in normally responding patients: a randomized trial. *Fertil Steril* 2011;95:812-4.
10. Karande VC, Morris R, Chapman C, Rinehart J, Gleicher N. Impact of the "physician factor" on pregnancy rates in a large assisted reproductive technology program: do too many cooks spoil the broth? *Fertil Steril* 1999;71:1001-9.
11. Maxwell KN, Cholst IN, Rosenwaks Z. The incidence of both serious and minor complications in young women undergoing oocyte donation. *Fertil Steril* 2008;90:2165-71.
12. Mohammed A, Woad KJ, Mann GE, Craigon J, Raine-Fenning N, Robinson RS. Evaluation of progestogen supplementation for luteal phase support in fresh in vitro fertilization cycles. *Fertil Steril* 2019;112:491-502.
13. Oliveira JB, Martins AM, Baruffi RL, Mauri AL, Petersen CG, Felipe V, et al. Increased implantation and pregnancy rates obtained by placing the tip of the transfer catheter in the central area of the endometrial cavity. *Reprod Biomed Online* 2004;9:435-41.
14. Purcell KJ, Schembri M, Telles TL, Fujimoto VY, Cedars MI. Bed rest after embryo transfer: a randomized controlled trial. *Fertil Steril* 2007;87:1322-6.
15. Tureck RW, Garcia CR, Blasco L, Mastroianni L Jr. Perioperative complications arising after transvaginal oocyte retrieval. *Obstet Gynecol* 1993;81:590-3.
16. van Os HC, Roozenburg BJ, Janssen-Caspers HA, Leerentveld RA, Scholtes MC, Zeilmaker GH, et al. Vaginal disinfection with povidone iodine and the outcome of in-vitro fertilization. *Hum Reprod* 1992;7:349-50.
17. van Weering HG, Schats R, McDonnell J, Hompes PG. Ongoing pregnancy rates in in vitro fertilization are not dependent on the physician performing the embryo transfer. *Fertil Steril*

2005:83:316-20.